



Original Article

Preliminary Phytochemical Screening, Quantification of phenolic compounds, of Plant Extract from *Chenopodium quinoa*

KADRI mounira ^{a*} RAMDANE Farah ^{b*} BOUZID Rachida^{c*} ATTALI Halima^c

^a University Elchahid Hamma lakhder. El-oued, Algeria and Biology, Environment and Health Laboratory, faculty of Nature Sciences and Life, Eloued, Algeria.

^b Faculty of Natural and Life Sciences, El Oued University BP 789, 39000, Algeria Biogeochemistry Laboratory in Desert Environments. Kasdi Merbah University. BP 511, 30000. Ouargla. Algeria

^c University Elchahid Hamma lakhder. El-oued, Algeria

ARTICLE INFOR

Article history:

Received 08 Marsh 2021

Revised 25 Mai 2021

Accepted 17 Jun 2021

Keywords:

Chenopodium quinoa

Polyphenol

Secondary metabolites

ABSTRACT

The aim of this study was to screening the phytochemical compounds of the seeds of *Chenopodium quinoa*, obtained from Wilaya of El-Oued.

The chemical study showed that the plant contained a number of secondary metabolites, flavonoids, tannins, saponins, steroids and triterpenes, glycosids while the absence of alkaloids and coumarin in *Chenopodium quinoa*.

Folin-Ciocalteu colorimetric were used to determine the total phenolic con-tent (TPC), in the hydroalcoholic seeds extracts. The yield of the methanolic extract was estimated at 36.66%. As for the quantitative content of polyphenols, it is $11.647 \pm 1.91 \mu\text{g AGE} / \text{mg extract}$

From this, *Chenopodium quinoa* is considered a nutritional and therapeutic value because it contains secondary metabolites

Faculty of Natural Sciences and Life, University of El Oued

1. Introduction

Coeliac disease is an immune-mediated enteropathy against dietary gluten present in wheat, rye and barley and is one of the most common lifelong food-related disorders worldwide [1]. in addition to the world 925 to 1023 million people were suffering chronic hunger [2]. One way of addressing solutions for this tremendous issue, is to look for crops that are nutritionally rich and can also grow under stressful environmental and research climatic conditions typical of the world hunger areas, such as Africa, among which species of the genus *Amaranthus* and *Chenopodium* [3]

Quinoa (*Chenopodium quinoa* Willd.) is a crop used by pre-Columbian cultures in South America for centuries. There is a long history of safe use of the grain in South America. Cultivated and collected *Chenopodium* species have been part of the Tiahuanacotan and Incan cultures. Quinoa has fulfilled various roles in these ancestral cultures, in addition to its role in human and animal nutrition, quinoa had a sacred importance [4].

Quinoa is usually referred to as a pseudo-cereal since it is not a member of the Gramineae family, but it produces seeds that can be milled in to flour and used as a cereal crop. it is considered to have a high nutritional value, mainly due to high in quantity and quality of protein relative to other protein sources [5]. Quinoa grains also have vitamins C, E (tocopherols), and B complex, and important minerals (Ca, K, Fe, Mg, Mn, P), isoflavons and even the saponins in the seed coats, previously considered as antinutrients [6].

2. Materials and Methods

2.1. Plant material

Quinoa (*Chenopodium quinoa*) was collected during Mars 2019, from Oum Thiour wilaya El M'Ghair is located within the Sahara Desert in northern-central of south Algeria.

* Corresponding author : KADRI Mounira E-mail address: mounira-kadri@univ-eloued.dz

Peer review under responsibility of University of El Oued

DOI : <http://doi.org/10.57056/ajb.v2i1.40>

2.2. Preparation of extracts

Total methanol extract of *Chenopodium quinoa* was prepared by maceration technique, the dried and powder of seeds plant (5g) were macerated with (20 ml) of methanol (70%) at room temperature 3 time (24 hours ×3). After filtration, the extract was concentrated using a rotary evaporator at a maximum temperature of 45°C, the residuals obtained were divided, a half part was stored in a freezer at – 4°C until further study [7].

2.3. extraction yield

The extraction yield is calculated by the formula given by [8]

$$R (\%) = 100 M / M'$$

R: is the yield in%;

M: the mass of the extract after evaporation of the solvent in mg

M': is the dry mass of the plant sample in mg.

2.4. Phytochemical screening

The methanol extracts were subjected to phytochemical tests for secondary metabolites, tannins, saponins, steroid, alkaloids and glycosides in accordance with [9] and [10] with little modification.

Tests for Flavonoids

Pieces of magnesium ribbon and 1 ml HCl concentrated were mixed with 5 ml methanolic extract after few minutes and pink or orange color showed the presence of flavonoid. [11].

Test for Saponins

5 ml of distilled water was mixed with methanolic extract in a test tube and it was mixed vigorously. The frothing was mixed, the foam appearance showed the presence of saponins.

Test for reducing sugars (Fehling's test)

The methanol extract (2ml) was added to boiling Fehling's solution (A and B) in a test tube. Obtaining a brick-red precipitate indicates the presence of the reducing compounds [12].

Test for tannins

To 2ml of methanolic solution of each extract, 2 drops of ferric chloride (FeCl₃ solution diluted to 1%) were added [13]. The appearance of a dark green color indicates the presence of catechic tannins. The appearance of a blue-

green color indicates the presence of gallic tannins

Test for Sterols and triterpenes

Sterols and terpenes were sought after by the Liebermann reaction. 5ml of the methanolic extract were evaporated. The residue is dissolved hot in 1 ml acetic anhydride and 1 ml of chloroform; we added 1 ml of concentrated sulfuric acid. The appearance, at the interphase, purple ring, turning blue then green, said a positive reaction.

Test for Alkaloids

2 ml of the methanolic extract, were treated separately with both reagents (Maeyer's, Hager's and Dragendorff's), after which it was observed whether the alkaloids were present or absent in the turbidity or precipitate formation. The appearance orange, brown or white precipitate indicates the presence of Alkaloids Respectively.

Test for coumarins

NaOH test - 2ml of extract was treated with 3ml of 10% sodium hydroxide in a test tube. If the solution turns to yellow color, then it contains coumarins.

2.5. Determination of polyphenols using the Folin-Ciocalteu reagent

Principle: The reagent consists of a mixture of phosphotungstic acid (H₃PW₁₂O₄₀) and phosphomolybdic acid (H₃PMo₁₂O₄₀). It is reduced during oxidation phenols, in a mixture of blue oxides of tungsten and molybdenum [14]. The color produced, whose maximum absorption is between 725 and 760 nm is proportional to the number of polyphenols present in the plant extracts. [15]

Implementation of the dosage

The control polyphenol, in general gallic acid. To perform the assay, 125µL of Folin-Ciocalteu reagent are added to 125µL of diluted extract (1mg.mL⁻¹) or point of range. After 3 min added to 1250 µL of Na₂CO₃ (75g.L⁻¹) are then added and 1ml water. After 90mn in the dark, the absorbance reading at 760 nm is done using a spectrophotometer. Through the standard range, calculates the average concentration of polyphenols present in the extract's plants in µg gallic acid equivalent per mg of extracts (µg AG E/mg extracts) [16];[17].

Statistical analysis

Averages were calculated using Excel 2016

3. Results and Discussion

The preparation of extracts from the seeds of *Chenopodium quinoa* was carried out by maceration methods at room temperature, was obtained with a yield of 36,66% (w/w) (Table1)

Table 1. extract methanolic Yields of *Chenopodium quinoa*

Extract	Color	Aspect	Yield%
MES	Brown	hygroscopic	33,66%

MES : methanolic extract of seeds Quinoa

Phytochemical screening

Investigations on the phytochemical screening of *Chenopodium quinoa* seeds extracts revealed the presence of flavonoids, tannins, saponins, steroids and triterpenes, glycosids while the absence of alkaloids and comarin in *Chenopodium quinoa* (Table 2). These compounds are known to be biologically active. Our results are in close agreement with that reported by (18; 19; 20) .

Determination of polyphenols

In recent years, the consumption of polyphenols has been increasing, largely due to its beneficial effects on health.

Through the standard range, calculates the average concentration of polyphenols present in the extracts plants. see Fig. 1. We described here the presence of high polyphenol content. As for the quantitative content of polyphenols, it is $11.647 \pm 1.91 \mu\text{g AGE} / \text{mg extract}$




Fig 1. Calibration curve for total polyphenols.

Table 2. Phytochemicals found in methanolic extract of <i>Chenopodium quinoa</i>	
Phytochemicals	Seeds
Flavonoids	+ -
Saponin	+++
Steroids	+++
Reducing sugars	++
Tannins	++
Alkaloids-wagner's-reagents	-
Alkaloids-Draghandroff's-reagents	-
Alkaloids-Mayer's-reagents	-
Coumarins	-
+++ : Strong intensity reaction, ++ : Medium intensity reaction + : Weak intensity reaction, - : Non detected + - : trace	

Conclusion

Quinoa is considered to be of nutritional and therapeutic value. There are several products derived from quinoa: puffed grains, flour, pasta, etc. However, more sophisticated products or those whose production requires the use of more advanced technologies are about to be exploited, such as the extraction of favonoid, saponin, starch and proteins, from quinoa milk, These products are considered as the economic potential of quinoa because they enhance not only nutritional, but also physicochemical characteristics, which go far beyond the food industry as they supply products to the chemical, pharmaceutical and cosmetic industries.

Acknowledgements

We are thankful to the researchers from the Laboratory of Environment and Health Laboratory, faculty of Nature Sciences and Life , Eloued, Algeria, for their support during this work.

References

1. Lindfors K, Ciacci C, Kurppa K, Lundin Knut E. A, Makharia Govind K, Luisa Mearin M, Murray Joseph A, Verdu Elena F. and Kaukinen Katri. Coeliac disease. *Nature reviews Disease Primers*. 2019, 5(1):3
DOI: 10.1038/s41572-018-0054-z
2. FAO. Quinoa: An ancient crop to contribute to world food security. Latin America and the Caribbean, 2011, pp: 3-14.
3. Capriles V D, Coelho K D, Guerra-Matías A C, and Arêas J A G. Effects of processing methods on amaranth starch digestibility and predicted glycemic Index. *Journal of Food Science*. 2008 , 73:H160-H164.
4. Bonifacio A. *Chenopodium* Sp.: Genetic resources, ethnobotany, and geographic distribution. *Food Reviews International*. 2003, 19: 1–7.
5. Avila Ruiz G. Exploring novel food proteins and processing technologies A case study on quinoa protein and high pressure high temperature processing. *Thesis submitted in fulfillment of the requirements for the degree of doctor. Wageningen University. Wageningen*. 2016 , p: 09.
6. Stikić R, Jovanović Z, Marjanović M, Đorđević S. The Effect of Drought on Water Regime and Growth of Quinoa (*Chenopodium quinoa* Willd.). *Research Grants: TR-31006 and EU FP7 project AREA. Serbia*. 2015, 2: 52-8000
7. Harrar A, activités antioxydante et antimicrobienne d'extraits de *Rhamnus alaternus* L. *Mémoire magister, Université Farhat Abbas (Sétif)*. 2012, 95p.
8. Falleh H, Ksouri R, Chaieb K, Karray-Bourauoui N, Trabelsi N, Boulaaba M, Abdelly C. Phenolic composition of *Cynara cardunculus* L. organs, and their biological activities. *C. R. Biologies*; 2008 , 331(5): 372-379 doi: 10.1016/j.crv.2008.02.008
9. Harborne J B. Phytochemical methods. A guide to modern techniques of plants analysis. *Elsevier Science*. 1998 , 3 :11.
10. Rahman G, Syed U Jan, Syed F, Samiullah S, Nusrat J. Preliminary Phytochemical Screening, Quantitative Analysis of Alkaloids, and Antioxidant Activity of Crude Plant Extracts from Ephedra intermedia Indigenous to Balochistan; *Hindawi e Scientific World Journal*. 2017 , 1-7. <https://doi.org/10.1155/2017/5873648>
11. Katiki L M, Chagas A C S, Takahira R K, Juliani H R, Ferreira J F S, Amarante A F T. Evaluation of *Cymbopogon schoenanthus* Essential Oil in Lambs Experimentally Infected With *Haemonchus contortus*. *Veterinary parasitology*. 2012 , 186(4): 312- 318. <https://doi.org/10.1016/J.Vetpar.2011.12.003>
12. Ayoola G A, Coker H A B, Adesegun S A, Adepoju-Bello A A, Obaweya K, EC Ezennia E C, Atangbayila T O. Phytochemical Screening and Antioxidant Activities of Some Selected Medicinal Plants Used for Malaria, Therapy in Southwestern Nigeria. *Tropical Journal of Pharmaceutical Research*. 2008 , 7 (3) : 1019-1024.
13. Kamal T, Ahmad M, Raji AA, Muayad S, Rahma , Muhammad N O, Preliminary phytochemical screening test of *Garcinia griffithii* Plant. *Innova Ciencia*. 2012 , 4(4): 68-74.
14. Ribéreau-Gayon P . Les composés phénoliques des végétaux. *Editions Dunod, Paris*. 1968 , 254 pp
15. López-Fernández O, Domínguez R, Pateiro M, Munekata Paulo E.S., Rocchetti G, and Lorenzo José M. Determination of Polyphenols Using Liquid Chromatography–Tandem Mass Spectrometry Technique (LC–MS/MS): A Review *Antioxidants (Basel)*. 2020 , 9(6): 479. doi: 10.3390/antiox9060479
16. Singleton V L, Orthofer R, Lamuela-Raventos R M. Analysis of Total Phenols and Other Oxidation Substrates and Antioxidants By Means of Folin- Ciocalteu Reagent Methode. *Enzymologie* . 1999 , 299: 152-178 [Http://Dx.Doi.Org/10.1016/S0076-6879\(99\)99017-1](http://dx.doi.org/10.1016/S0076-6879(99)99017-1).
17. Slinkard K, Singleton V L. Total Phenol Analys: Automation and Comparison With manual Methods. *American journal of enology and viticulture*. 1977 , 28: 49-55.
18. Gordillo-Bastidas E, Díaz-Rizzolo DA, Roura E, Massanés T, Gomis R. - Quinoa (*Chenopodium quinoa* Willd). From Nutritional Value to Potential Health Benefits: An Integrative Review. *J Nutr Food Sci*. 2016, 6(3): 2155-9600 DOI: 10.4172/2155-9600.1000497
19. Jancurová M, Mínarovičová L, Dandár A, - Quinoa – a review. *Czech J. Food Sci*. 2009, Vol. 27: 71–79.
20. Maradini A, Pirozi M, Borges J, Sant'Ana H, Chaves J, Coimbra J, - Quinoa: Nutritional, functional and antinutritional aspects. *Crit Rev Food Sci Nutr*. 2017 , 24;57(8):1618-1630. doi: 10.1080/10408398.2014.1001811.

Recommended Citation

KADRI M., RAMDANE F., BOUZID R & ATTALI H. Preliminary Phytochemical Screening, Quantification of phenolic compounds, of Plant Extract from *Chenopodium quinoa*. *Alger. j. biosciences*. 2021, 02(01):042-045.



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/)