



## Original Article

# Measurement of radioactivity levels in powdered milk consumed in Algeria and estimation of annual effective doses

Linda Chabouni\*, Nesrine Hamitouche, Raouen Graichi, Ali Benbourenane, Omar Amri

Radiological expertise department, Nuclear Research Center of Algiers, 02, Bd Frantz Fanon B.P. 399 Alger-Gare, Algeria

### ARTICLE INFO

#### Article history:

Received 10 July 2023

Revised 08 November 2023

Accepted 11 November 2023

#### Keywords:

Radioactivity;  
Radiation dose;  
Radionuclides;  
Powdered milk; Gamma  
spectrometry;  
HPGe detector.

### ABSTRACT

The monitoring of radioactivity in food, especially in milk, has become a top priority in many countries. This paper presents the activity concentrations of  $^{40}\text{K}$ ,  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$  and  $^{137}\text{Cs}$  radionuclides measured in twenty-two samples of the powdered milk available in the Algerian market, sourced from different regions worldwide. The measurements were conducted using an HPGe detector by employing the gamma spectrometry method. The results show that the specific activity of  $^{40}\text{K}$ ,  $^{226}\text{Ra}$  and  $^{232}\text{Th}$  in infant powdered milk samples varied from  $(133.57 \pm 1.75$  to  $195.95 \pm 2.56$  Bq/kg),  $(1.35 \pm 0.03$  to  $2.70 \pm 0.07$  Bq/kg) and  $(1.34 \pm 0.03$  to  $1.63 \pm 0.06$  Bq/kg), respectively. In adult powdered milk samples, the activity concentration of  $^{40}\text{K}$ ,  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$  and  $^{137}\text{Cs}$  were varied from  $(110.52 \pm 1.49$  to  $687.89 \pm 8.94$  Bq/kg),  $(1.38 \pm 0.04$  to  $2.59 \pm 0.10$  Bq/kg),  $(1.25 \pm 0.03$  to  $2.63 \pm 0.09$  Bq/kg) and  $(3.60 \pm 0.07$  to  $7.78 \pm 0.11$  Bq/kg), respectively. The obtained results were subsequently compared with data from various studies conducted globally. The estimated annual ingestion dose resulting from the consumption of powdered milk was found to be  $583.80 \mu\text{Sv/y}$  infants ( $\leq 1\text{y}$ ), and  $56.85 \mu\text{Sv/y}$  for adults ( $\geq 17\text{y}$ ). These values indicate that powdered milk in Algeria does not have a significant radiological impact on the population, as they are below the dose limit recommended by the World Health Organization.

## 1. Introduction

Monitoring radioactivity has become a common practice in many countries, especially regarding food products and the environment. This approach plays a crucial role in safeguarding public health by regulating human exposure to radiation [1].

The primary sources of internal radiation exposure in humans through food ingestion are the naturally occurring radionuclides, particularly  $^{40}\text{K}$ , and the radionuclides from the  $^{238}\text{U}$  and  $^{232}\text{Th}$  series, which are extensively present in the environment [2]. In addition to natural radionuclides, various artificial radioactive elements, such as  $^{137}\text{Cs}$ , have the ability to spread through the food chain via food consumption. These elements have been introduced into our environment as a result of numerous nuclear weapons

experiments and various incidents involving nuclear reactors [3,4]. One of the primary food items that necessitate examination for its radioactivity levels is milk. It is an essential component of human dietary needs, particularly for infants under 1-year-old and children. Furthermore, milk acts as a vector of radioactive contamination in the food chain since it can be contaminated by radionuclides like  $^{137}\text{Cs}$ . These radionuclides can be transferred to milk from the grass ingested by the cows, which is laden with radioactive isotopes [5].

In Algeria, although milk is considered as a fundamental component of the diet, a considerable portion, especially powdered milk, is imported, primarily from Europe. That's

\*Corresponding author.

E-mail address: [l.chabouni@crna.dz](mailto:l.chabouni@crna.dz)

Peer review under responsibility of University of El Oued.

2716-9227/© 2023 The Authors. Published by University of El Oued. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>). <https://dx.doi.org/10.57056/ajet.v8i2.131>

why Algerian regulations require strict monitoring of radioactive contamination in this product.

The purpose of this study is to perform a qualitative and quantitative analysis of the radionuclides found in powdered milk consumed in Algeria, and subsequently estimate the annual ingestion dose for various age groups in the population resulting from this consumption.

## Materials and Methods

### 2.1. Samples conditioning

Twenty-two samples of powdered milk, imported from various countries and consumed in Algeria, were chosen for analysis. The origins and types of the samples are detailed in Table 1. A quantity of 250 g of each sample was packed in a marinelli beaker (Fig 1), and sealed for a

duration of four weeks to achieve the state of radioactivity equilibrium between the parent and daughter radionuclides.



Fig 1. Samples conditioning in Marinelli Beakers

Table 1. Sample codes and countries of origin for analyzed powdered milk samples.

	Sample code	Origin	Sample code	Origin	
Adult milk powder	Sample 01	Netherlands	Sample 14	France	
	Sample 02	Ireland	Sample 15	Denmark	
	Sample 03	Argentina	Infant milk powder	Sample 16	France
	Sample 04	Holland		Sample 17	Germany
	Sample 05	Ukraine		Sample 18	Switzerland
	Sample 06	Lithuania		Sample 19	France
	Sample 07	Belgium		Sample 20	France
	Sample 08	Poland		Sample 21	Switzerland
	Sample 09	Canada		Sample 22	France
	Sample 10	France			
Sample 11	New Zealand				
Sample 12	Uruguay				
Sample 13	USA				

### 2.2. Gamma Spectrometric Analysis

The powdered milk samples were analyzed using a high-resolution gamma-ray spectrometry system (Fig 2) that incorporated a coaxial hyper-pure germanium detector (Canberra model GC3018). This detector has an energy resolution of 1.8 keV at the 1.33 MeV  $\gamma$ -ray line of  $^{60}\text{Co}$  and an efficiency of 30%. The system also included a preamplifier (model 2002CSL) positioned in close proximity to the detector, an amplifier (model 2024A), and a 16 k digital multichannel analyzer. In order to maintain a low background environment, the detector was enclosed within a 10 cm steel shield coated internally with a 1 mm copper layer.

The energy calibration is performed using a multi-nuclides gamma-ray reference source ( $^{241}\text{Am}$ ,  $^{137}\text{Cs}$  and  $^{60}\text{Co}$ ) that covered a wide energy range from 59.5 to 1332.5 keV, while the efficiency calibration was done using a standard source that has the same matrix and same geometry as the measuring samples; it is powder milk contaminated with a

radioactive source of  $^{152}\text{Eu}$ , with an activity of 326 Bq in 1993, emitting gamma rays in the energy range of 121–1408 keV.



Fig 2. Gamma spectrometry system

By employing this standard source, the counting efficiency curve of the HPGe detector was drawn and is calculated using the following equation:

$$\xi = \frac{N}{A * I\gamma * tc} \quad (1)$$

Where N is the net area at the interest peak energy, corrected for background counts, A represents the source activity in Becquerel with decay time correction; tc stands for the collection time in seconds and  $I\gamma$  signifies the intensity of gamma ray.

Figure 3 and 4 show respectively the energy calibration line and the curve of the efficiency calibration.




Fig 3. Energy calibration line.




Fig 4. Efficiency calibration curve.

The samples were placed above the detector for 14400 s to minimize statistical counting errors. The background was also measured for the empty container with the same counting time and subtracted from the samples spectra. The figure below illustrates an example of spectra from an adult milk powder sample obtained using the Genie 2000 software.




Fig 5. Gamma ray spectrum of Adult milk Powder sample (Sample 05: Ukraine).

**2.3. Activity calculation:**

The activity concentration of radionuclides in milk samples, expressed in Bq/kg, was determined using the following equation [6]:

$$A = \frac{N}{\xi * I\gamma * tc * m} \quad (2)$$

The uncertainty related to the activity concentration is determined through error propagation, as described by the following relationship [7]:

$$\delta A = A \sqrt{\left(\frac{\sigma_{\xi}}{\xi}\right)^2 + \left(\frac{\sigma_N}{N}\right)^2 + \left(\frac{\sigma_{I\gamma}}{I\gamma}\right)^2 + \left(\frac{\sigma_m}{m}\right)^2 + \left(\frac{\sigma_{tc}}{tc}\right)^2} \quad (3)$$

Where N represents the net peak counts after background subtraction;  $\xi$ , corresponds to the detector counting efficiency at the specific gamma ray energy; tc, is the measured time in seconds; I, represents the intensity of  $\gamma$ -ray; while m indicates the weight of the sample in kilograms.

The activity concentration of  $^{226}\text{Ra}$  and  $^{232}\text{Th}$  was assessed by relying on their short-lived progenies, utilizing their distinct and notably strong gamma spectral lines [10]. This approach was necessary due to the limited intensity of gamma rays emitted by  $^{226}\text{Ra}$  and  $^{232}\text{Th}$ , making them undetectable by conventional gamma-ray spectrometry [8,9].

The activity concentrations of  $^{226}\text{Ra}$  and  $^{232}\text{Th}$  were then calculated using equations 4 and 5 respectively:

$$A_{226\text{Ra}}(\text{Bq / kg}) = \frac{A_{214\text{Bi}} + A_{214\text{Pb}}}{2} \quad (4)$$

$$A_{232\text{Th}}(\text{Bq / kg}) = \frac{A_{228\text{Ac}} + A_{212\text{Pb}} + A_{208\text{Tl}}}{3} \quad (5)$$

On the other hand, the activity concentrations of  $^{40}\text{K}$  and  $^{137}\text{Cs}$  were directly assessed using their individual gamma lines at 1460.8 keV and 661.6 keV, respectively.

#### 2.4. Estimation of annual effective dose

To calculate the annual effective dose (D) resulting from the ingestion of radionuclides through the consumption of powdered milk, we used the following equation [11]:

$$D = AIE \quad (6)$$

Where D is the annual effective dose in (Sv/y), A is the activity concentrations of the interested radionuclides in the sample (Bq/kg), I is the annual powdered milk intake (kg/y), which depends on the human age (for infants  $\leq 1$  year is 22.4 kg/y and adult  $>17$  year is 13 kg/y) [12] and E is the dose conversion factors for the radionuclides (Sv/Bq). The conversion factor E varies depending on both the radioisotopes and the ages of the individuals (Table 2).

Table 2: Dose conversion factors of radionuclides for infants ( $\leq 1$ ) and adults ( $>17$  y) [13].

Age group	Dose conversion factors (Sv/Bq)			
	$^{40}\text{K}$	$^{226}\text{Ra}$	$^{232}\text{Th}$	$^{137}\text{Cs}$
Infants ( $\leq 1$ y)	$6.2 \times 10^{-8}$	$4.7 \times 10^{-6}$	$4.6 \times 10^{-6}$	$2.1 \times 10^{-8}$
Adults ( $> 17$ y)	$6.2 \times 10^{-9}$	$2.8 \times 10^{-7}$	$2.3 \times 10^{-7}$	$1.3 \times 10^{-8}$

The annual effective dose resulting from the consumption of radionuclides found in powdered milk are tabulated in Table 5.

## 2. Results and Discussion

### 3.1. Activity Concentration:

The activity concentrations of radionuclides measured in the powdered milk samples are given in Table 3. The table

indicates that the activity concentrations of  $^{40}\text{K}$  in all samples were higher than for  $^{226}\text{Ra}$  and  $^{232}\text{Th}$ . This can be attributed to the natural abundance of potassium-40 in the environment. Its widespread presence in the Earth's crust and soil allows it to be easily incorporated into plants and subsequently transferred through the food chain, making it an integral part of the biological processes within ecosystems [14].

The measurement results also reveal that the specific activities of  $^{40}\text{K}$ ,  $^{226}\text{Ra}$ , and  $^{232}\text{Th}$  exhibit variation across different samples. In infant powdered milk samples, the activity concentrations varied from  $133.57 \pm 1.75$  (Germany) to  $195.95 \pm 2.56$  Bq/kg (France) for  $^{40}\text{K}$ , from  $1.35 \pm 0.03$  to  $2.70 \pm 0.07$  Bq/kg for  $^{226}\text{Ra}$ , and from  $1.34 \pm 0.03$  to  $1.63 \pm 0.06$  Bq/kg for  $^{232}\text{Th}$ , respectively.

In adult powdered milk samples, the activity concentrations of  $^{40}\text{K}$ ,  $^{226}\text{Ra}$ , and  $^{232}\text{Th}$  varied from  $110.52 \pm 1.49$  (New Zealand) to  $687.89 \pm 8.94$  Bq/kg (USA), from  $1.38 \pm 0.04$  to  $2.59 \pm 0.10$  Bq/kg, and from  $1.25 \pm 0.03$  to  $2.63 \pm 0.09$  Bq/kg, respectively.

This variation can be attributed to diverse factors including background levels of natural radioactivity in various regions of the world, the transfer of radionuclides from soil to plants/grass, the geological characteristics of local dairy farming areas, climate and agricultural practices in various farming areas, the intake of feedstuffs by different animals (cow in this case), and the raw materials involved in processing [15].

Besides natural radioactivity, the presence of artificial radioactivity ( $^{137}\text{Cs}$ ) was also detected in two samples of adult milk powder from Poland ( $3.60 \pm 0.07$  Bq/kg) and Ukraine ( $7.78 \pm 0.11$  Bq/kg). However, the measured activities are below regulatory limits [16]. The occurrence of this radionuclide in powdered milk is a result of its dispersion in the environment due to nuclear activities or accidents [17].

Table 3: Activity concentrations of different radionuclides found in milk powder samples.

Samples code	Activity concentration (Bq/kg)			
	$^{40}\text{K}$	$^{226}\text{Ra}$	$^{232}\text{Th}$	$^{137}\text{Cs}$
Sample 01	$552.28 \pm 7.18$	$2.59 \pm 0.10$	$2.63 \pm 0.09$	BDL
Sample 02	$685.18 \pm 8.90$	$2.46 \pm 0.09$	$1.63 \pm 0.05$	BDL
Sample 03	$466.84 \pm 6.07$	$1.38 \pm 0.04$	$1.46 \pm 0.04$	BDL
Sample 04	$568.21 \pm 7.38$	$2.26 \pm 0.07$	$1.71 \pm 0.06$	BDL
Sample 05	$558.72 \pm 7.26$	$2.56 \pm 0.08$	$2.04 \pm 0.08$	$7.78 \pm 0.11$

Sample 06	597.37±7.76	1.93±0.07	1.49±0.05	BDL
Sample 07	595.67±7.74	2.32±0.08	1.91±0.07	BDL
Sample 08	619.40±8.05	1.74±0.06	1.50±0.04	3.60±0.07
Sample 09	632.29±8.22	2.34±0.09	1.57±0.05	BDL
Sample 10	668.90±8.69	2.13±0.07	1.79±0.06	BDL
Sample 11	110.52±1.49	1.62±0.06	1.25±0.03	BDL
Sample 12	383.10±4.98	1.78±0.05	1.56±0.04	BDL
Sample 13	687.89±8.94	1.83±0.08	1.43±0.03	BDL
Sample 14	179.00±2.34	1.75±0.06	1.52±0.05	BDL
Sample 15	165.10±2.16	1.29±0.04	1.37±0.04	BDL
Sample 16	182.05±2.38	2.46±0.08	1.63±0.06	BDL
Sample 17	133.57±1.75	2.97±0.09	1.51±0.04	BDL
Sample 18	171.88±2.28	1.35±0.03	1.44±0.05	BDL
Sample 19	174.93±2.29	1.83±0.07	1.47±0.06	BDL
Sample 20	141.37±1.85	1.49±0.05	1.46±0.04	BDL
Sample 21	144.08±1.89	1.71±0.06	1.34±0.03	BDL
Sample 22	195.95±2.56	2.70±0.07	1.50±0.05	BDL

BDL: below detection limit.

Table 4: Comparison of activity concentration of radionuclides detected in powdered milk with worldwide values.

Country	Activity Concentration (Bq/kg)				Reference
	<sup>226</sup> Ra	<sup>232</sup> Th	<sup>40</sup> K	<sup>137</sup> Cs	
Thailand <sup>I</sup>	3.20±0.51	1.24±0.10	231.20±25.50	4.90±0.74	[18]
Saudia Arabia <sup>I</sup>	0.46±0.05	0.35±0.08	234.18±1.90	N/A	[19]
Malaysia <sup>I</sup>	3.05±0.45	2.55±0.64	99.10±5.58	0.27±0.19	[20]
Bangladesh <sup>I</sup>	5.40±0.40	5.70±0.50	334.40±12.00	N/A	[21]
Iraq <sup>I</sup>	3.88±0.56	1.58 ±0.39	387.18±6.29	N/A	[22]
Egypt <sup>I</sup>	0.91 ± 0.20	0.60 ± 0.11	478,0 ± 25,00	0.42 ±0.11	[23]
Algeria <sup>I</sup>	N/A	N/A	106.70±5.44	N/A	[24]
Iraq <sup>A</sup>	BDL	BDL	290.67±14.97	N/A	[25]
Brazil <sup>A</sup>	BDL	0.50	489.52±13.00	2.54	[3]
Iran <sup>A</sup>	0.11±0.02	0.13±0.02	549.65±14.97	1.43±0.04	[26]
Jordan <sup>A</sup>	1.24±0.77	N/A	348.30±7.76	0.52±0.39	[5]
Malaysia <sup>A</sup>	1.97 ±0.32	1.07±0.24	307.74 ± 7.20	N/A	[27]
Saudia Arabia <sup>A</sup>	9.64±0.45	6.77±0.40	74.51±6.40	N/A	[28]
India <sup>A</sup>	2.50±1.20	N/A	34.35±5.20	N/A	[29]
Algeria <sup>I</sup>	1.95±0.06	1.47±0.05	165.5±2.20	BDL	Present work
Algeria <sup>A</sup>	2.07±0.07	1.69±0.05	548.17±7.12	5.69±0.09	Present work

BDL: below detection limit, N/A represents data not available, I: Infant milk powder, A: Adult milk powder.

The findings presented in this study were compared to the results obtained in other studies worldwide, which focused

on the activities of radionuclides in powdered milk. These comparative values are summarized in Table 4. As shown

in Table 4, our study found that the average activity concentrations of  $^{40}\text{K}$  in powdered milk for infants were higher than those in Malaysia but lower than levels detected in powdered milk from several other regions. Regarding adult samples, the average concentration was higher than what previous research had reported, except for Iran. Concerning the activity values of  $^{226}\text{Ra}$  and  $^{232}\text{Th}$ , as they are compared to published data from various countries, it is evident that the majority of these values are within the range of the published data. In the case of  $^{137}\text{Cs}$ , the average radioactivity concentrations in this study are relatively elevated when compared to levels found in other nations. The results indicate that the differences in the levels of radioactivity in milk can be explained by differences in their original surroundings and/or the raw materials used in their production [25].

### 3.2. The Annual Effective Dose for milk ingestion

Table 5 illustrate the annual effective dose calculations resulting from the ingestion of radionuclides in powdered milk for different age groups.

Table 5: Annual effective dose resulting from the ingestion of radionuclides.

Samples Code		Annual effective dose ( $\mu\text{Sv/y}$ )			
		$^{40}\text{K}$	$^{226}\text{Ra}$	$^{232}\text{Th}$	$^{137}\text{Cs}$
Powder milk (Adults >17Y)	1	44.51	9.44	7.86	BDL
	2	55.23	8.96	4.88	BDL
	3	37.63	5.03	4.36	BDL
	4	45.80	8.24	5.14	BDL
	5	45.80	9.34	6.12	1.3
	6	48.15	7.03	4.46	BDL
	7	48.01	8.46	5.72	BDL
	8	49.92	6.35	4.49	0.61
	9	50.96	8.53	4.69	BDL
	10	53.91	7.78	5.36	BDL
	11	8.91	5.90	3.75	BDL
	12	30.88	6.49	4.67	BDL
	13	55.44	6.69	4.28	BDL
	<b>Average</b>	<b>44.24</b>	<b>7.55</b>	<b>5.06</b>	<b>0.95</b>
Powder milk (Infants $\leq 1\text{Y}$ )	14	248,61	183,98	156,07	BDL
	15	229,30	135,63	140,72	BDL
	16	252,84	259,22	168,03	BDL
	17	185,51	312,69	155,57	BDL
	18	238,72	142,64	148,34	BDL
	19	242,96	169,01	151,08	BDL
	20	196,34	156,87	150,10	BDL
	21	200,11	180,10	139,10	BDL
	22	272,15	284,24	154,35	BDL
		<b>Average</b>	<b>248,61</b>	<b>183,98</b>	<b>156,07</b>

The results showed that  $^{40}\text{K}$  is the primary factor influencing the internal radiation dose experienced by the

population through the consumption of powdered milk. Moreover, the average annual effective dose resulting from  $^{40}\text{K}$  intake in infants is significantly higher in comparison to that in adults. This difference can be attributed to the fact that milk acts as the primary nutritional source for infants in their initial six months of life, leading to a higher rate of consumption [19]. In the case of  $^{137}\text{Cs}$ , the dose resulting from the consumption of this radionuclide is negligible, indicating that it does not present a potential health hazard. The findings of this study demonstrate that the annual effective doses resulting from the consumption of radionuclides in powdered milk are all below the recommended limit of 1 mSv/y established by the Joint Food and Agriculture Organization of the United Nations/World Health Organization (FAO/WHO) [30] and the International Commission on Radiological Protection (ICRP) [31].

## 4. Conclusion

The activity concentrations of radionuclides were measured in 22 samples of powdered milk imported from different countries and marketed in Algeria using High-resolution Gamma-ray spectrometry with H5PGe detector.

The results show that the average activity concentrations of  $^{40}\text{K}$ ,  $^{226}\text{Ra}$  and  $^{232}\text{Th}$  in infant powdered milk samples were  $165.32 \pm 2.16$ ,  $1.95 \pm 0.06$  and  $1.47 \pm 0.05$  Bq/kg, respectively.

In adult powdered milk samples, the activity concentration of  $^{40}\text{K}$ ,  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$  and  $^{137}\text{Cs}$  were  $548.18 \pm 7.12$ ,  $2.07 \pm 0.07$ ,  $1.69 \pm 0.05$  and  $5.69 \pm 0.09$  Bq/kg, respectively.

The annual effective dose resulting from the consumption of these powdered milk samples by infants and adults was also evaluated. The overall result indicates that the radionuclide levels in the studied samples of powdered milk for children and adults are clearly below the recommended limits issued by FAO/WHO. In general terms, it can be concluded that the powdered milk consumed in Algeria is radiologically secure and is unlikely to pose substantial health risks. However, it is advisable to maintain continuous monitoring of the radiological safety of milk consumed by the population in Algeria.

## Conflict of Interest

The authors declare that they have no conflict of interest

## References

1. Francesco C, Valentina V, Giuseppe P, et al. Assessment of Radioactivity Concentration in Milk Samples Consumed in Italy. *Current Nutrition & Food Science*, 2023; 19: 176-181.
2. Mohsin K. Study of Radioactivity of Selected Samples of Soil in Amarah City, Maysan Province, Iraq. *Engineering and Technology Journal*, 2018, 36 Part C.
3. Melquiades FL and Appoloni CR.  $^{40}\text{K}$ ,  $^{137}\text{C}$  and  $^{232}\text{Th}$  activities in Brazilian milk samples measured by gamma ray spectrometry. *Ind. J. Pure. Appl. Phys.* 2002; 40:5–11.
4. Mlwilo NA, Mohammed Najat K and Spyrou NM. Radioactivity levels of staple foodstuffs and dose estimates for most of the Tanzanian population. *Journal of radiological Protection*, 2007; 27: p. 471.
5. Ababneh ZQ, et al. Measurement of natural and artificial radioactivity in powdered milk consumed in Jordan and estimates of the corresponding annual effective dose. *Radiation protection dosimetry*, 2010; 138(3): 278-283.
6. Dovlete C, Povinec PP. Quantification of uncertainty in gamma-spectrometric analysis of environmental samples. *Quantifying uncertainty in nuclear analytical measurements*. 2004;103.
7. Génie2000, Spectrometry Software, Software Analysis Algorithms, Technical Documentation.
8. Dowdall M, Selnaes ØG, Gwynn JP, et al. Simultaneous determination of  $^{226}\text{Ra}$  and  $^{238}\text{U}$  in soil and environmental materials by gamma-spectrometry in the absence of radium progeny equilibrium. *Journal of radioanalytical and nuclear chemistry*. 2004; 261: 513-521.
9. El mestikou R, Jemii E, Mazouz M, et al. Determination of the activity level in powdered milk available in Tunisia and assessment of the radiological risks. *Journal of Radioanalytical and Nuclear Chemistry*, 2018; 317: 991-996.
10. Scholten Jan C, Osvath Iolanda Pham, Mai K.  $^{226}\text{Ra}$  measurements through gamma spectrometric counting of radon progenies: How significant is the loss of radon?. *Marine Chemistry*, 2013; 156: 146-152.
11. UNSCEAR. United Nations Scientific Committee on the Effects of Atomic Radiation. Sources and effects of ionizing radiation .Report to General Assembly (New York: United Nations) .2000.
12. UNSCEAR. United Nations Scientific Committee on the Effects of Atomic Radiation. Sources and effects of ionizing radiation. Report to General Assembly (New York: United Nations).1993.
13. Eckerman K. et al. ICRP Publication 119: Compendium of dose coefficients based on ICRP Publication 60, *Ann. ICRP*, 2012; 41: 1–130.
14. Uwatse O.B et al. Radiation dose to Malaysian infants from natural radionuclides via consumption of powdered milk. *AIP Conference Proceedings*. 2015; 1657.
15. Badran H M, Sharshar T, Elnimer T. Levels of  $^{137}\text{Cs}$  and  $^{40}\text{K}$  in edible parts of some vegetables consumed in Egypt. *Journal of Environmental Radioactivity*, 2003; 67: 181-190.
16. Presidential Decree 05/117 of 11 Apr. 2005 on measures to protect against ionizing radiation. O.J.R.A.2005, 27: 3-24.
17. CAI Junjie IP, Kwai Fun EZE, Chika et al. Dispersion of radionuclides released by nuclear accident and dose assessment in the Greater Bay Area of China. *Annals of Nuclear Energy*, 2019; 132: 593-602.
18. Poltabtim W, & Saenboonruang K. Assessment of activity concentrations and their associated radiological health risks in commercial infant formulas in Thailand. *Chiang Mai Journal of Science*, 2019;46(4): 778-786.
19. Al-zahrani JH. Natural radioactivity and heavy metals in milk consumed in Saudi Arabia and population dose rate estimates. *Life Science Journal*, 2012; 9 (2):651-656.
20. Uwatse OB, Olatunji MA, Khandaker MU, Amin YM, Bradley DA, Alkhorayef M, & Alzimami K. Measurement of natural and artificial radioactivity in infant powdered milk and estimation of the corresponding annual effective dose. *Environmental Engineering Science*, 2015; 32: 838-846.
21. Begam K., Rahman MM, Kabir MA, Tamim U, Hossain SM, & Begum A. Natural radioactivity level of  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  in baby food and committed annual effective dose assessment in Bangladesh. *International Journal of Environmental Monitoring and Analysis*, 2020; 8: 187.
22. Al-jnabi MKM. Estimating Radioactivity in Various Types of Milk using NaI (TI) detector. *journal of kerbala university*, 2017; 13: 229-236.
23. Salahel din K. Assessment of natural and artificial radioactivity in infants' powdered milk and their associated radiological health risks. *Journal of Radioanalytical and Nuclear Chemistry*, 2020; 324: 977-981.
24. Kasmi Y, Kouachi S. Radio-isotopic control of imported infant formula. Engineering Degree Thesis. *Higher School of Food Science and Agri-Food Industry*, 2021.
25. Sahar A, Rana AM, Al-ani R. Assessment of natural radionuclides in powdered milk consumed in Iraq. *Assessment*, 2016; 13.
26. Hosseini T, Fathivand AA, Barati H, Karimi M. Assessment of radionuclides in imported foodstuffs in Iran. *Iran J Radiat Res*.2006; 4(3):149–153.
27. Priharti W, Samat SB, Yasir MS. Assessment of radiation hazard indices arising from natural radionuclides content of powdered milk in Malaysia. *Journal of Radio analytical and Nuclear Chemistry*, 2016; 307: 297-303.
28. Alamoudi Zain M. Assessment of natural radionuclides in powdered milk consumed in Saudi Arabia and estimates of the corresponding annual effective dose. *Journal of American science*, 2013; 9: 267-273.

29. Shanthi G, kumaran J, Thampi Thanka RAJ G. Allan Gnana, et al. Natural radionuclides in the South Indian foods and their annual dose. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 2010; 619, 1-3: 436-440.
30. FAO/WHO, Codex alimentarius, general requirements, section 6.2, guideline levels for radionuclides in foods following accidental nuclear contamination for use in international trade, Joint FAO/ WHO Food Standards Programme, Rome.1995.
31. ICRP. The 2007 Recommendations of the International Commission on Radiological Protection. Publication 103, *Annals of the ICRP*, 37.2007.

### Recommended Citation

Chabouni L, Hamitouche N, Graichi R, Benbourenane A, Amri O. Measurement of radioactivity levels in powdered milk consumed in Algeria and estimation of annual effective doses, *Alger. J. Eng. Techno.* 2023; 8(2):258-265. <https://dx.doi.org/10.57056/ajet.v8i2.131>



This work is licensed under a [Creative Commons Attribution-Non Commercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/)